

Dequadin[®]

0.25mg Lozenges
DEQUALINIUM CHLORIDE

FOR SORE THROATS
AND MOUTH INFECTIONS

Reduces tenderness of the throat and mouth
Relieves symptoms of mouth ulcers and infections

20 Lozenges

Dequadin[®]

0.25mg Lozenges

FOR SORE THROATS
AND MOUTH INFECTIONS

Dequadin[®]

0.25mg Lozenges

For the relief of symptoms of sore throats and mouth infections including mouth ulcers, tonsillitis, stomatitis and oral thrush

Do not take:

- If you are under 10 years old
- If you are allergic to any of the ingredients
- If you have been told by your doctor that you have an intolerance to some sugars.

Talk to your pharmacist or doctor:

- if your symptoms do not go away after a few days
- if you take too many and feel unwell
- if you are concerned or think that you may be reacting badly to this medicine.

Check before use:

- Diabetics- take account of the carbohydrate content
- Consult your doctor before use if you are pregnant or breast feeding.

WARNING: DO NOT EXCEED THE STATED DOSE.

How to use

Read the carton carefully before use.

Adults & Children over 10 years: Suck one lozenge slowly every 2-3 hours. Do not take any more than 8 lozenges in any 24 hour period.

Side Effects

Soreness of the tongue and allergic reactions to the ingredients may occur.

What is in Dequadin 0.25mg Lozenges?

- Active Ingredient: Dequalinium Chloride 0.25mg.
- Also contains: Icing Sugar, Citric Acid Monohydrate, Liquid Glucose, Sodium Saccharin, Camphor, Magnesium Stearate, Gelatin, Sunset Yellow (E110), Flavours (Lime and Peppermint Oil).
- Each lozenge contains 1g of Sucrose (from Glucose and Fructose) and 50mg of Glucose.
- Sunset Yellow (E110) may cause allergic reactions.

KEEP ALL MEDICINES OUT OF REACH AND SIGHT OF CHILDREN.

Text Prepared: November 2011

