

# FRADOR

CHLORBUTOL 1.0% w/v

• Gives prompt relief from the pain associated with mouth ulcers and gum sores and on-going protection from saliva, bacteria and food whilst healing proceeds. Frador should be applied 4 times a day after meals and before going to bed. In the event of missing an application, apply as needed.

**DIRECTIONS:**

Stand bottle in tray provided. Dry Ulcer. Using applicator, apply Frador directly to mouth ulcer. There may be a temporary tingling sensation. Keep mouth open for several seconds to allow Frador to dry. Application is complete. If healing does not start within a few days always consult your doctor.

**CAUTION:**

As with all medicines - KEEP OUT OF REACH OF CHILDREN. For adult use only. Avoid use in pregnancy. Keep away from naked flame. Retain carton for instructions. Not to be taken internally. In rare occurrences users can experience a local reaction to some or all of the ingredients. If excess irritation occurs during or after application, stop use. If condition persists, contact your pharmacist or doctor. There have been no reports of ill-effects from overdosage. Store at room temperature. Avoid spilling on polished surfaces or fabrics. Spilt Frador can be removed from hands and fingers with methylated spirits.

**REPORTING OF SIDE EFFECTS:**

If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via the Yellow Card Scheme at: [www.mhra.gov.uk/yellowcard](http://www.mhra.gov.uk/yellowcard). By reporting side effects you can help provide more information on the safety of this medicine.

Licence Holder: Castagnola Ltd, London NW3 6AL

Date of Revision of Text: 19/06/2015

Contract Manufacture: Thorpe Laboratories Ltd, LN12 1NB, Distributor: Fenton Pharmaceuticals Ltd, London NW3 6AL

PL 01520/0001

G.S.T

Batch No:  
Empty Date:

Active Ingredients: Chlorbutol B.P. 1.0% w/v  
Eulemnic Benzoin 1.0% w/v, Prop Stryx B.P.C. 2.8% w/v,  
Eulemnic Benzoin 1.0% w/v, in Alcohol Solution to 100% Carminols.

# FRADOR



*For mouth ulcers and gum sores*

3.5ml

- Immediate effective relief
- Soothes pain & aids healing
- Hygienic application

CHLORBUTOL 1.0% w/v

# FRADOR

**FRADOR**  
For mouth ulcers