

The safety of Medijel Gel during pregnancy or lactation has not been established. Pregnant and breast-feeding mothers should speak to a doctor or pharmacist before using product. Contains small amounts of ethanol (alcohol), less than 100mg per dose. If you have been told by your doctor that you have an intolerance to some sugars, contact your doctor before taking Medijel Gel.

Contains Sucrose. Oral Gel **Active Ingredients:** Lidocaine Hydrochloride 0.66% w/w, Aminoacridine Hydrochloride 0.05% w/w.

Also contains Sucrose, Ethanol (96%), Carbomer, Diisopropanolamine 90% Aqueous, Hydroxypolyethoxydodecane, Peppermint Oil, Ethyl Vanillin, Saccharin Sodium, Glycerol, Purified Water.

Do not use after the expiry date stated on the pack.

Keep all medicines out of the reach and sight of children.

PL 0133/5000R MA 137/00301

Made by the MA holder DDD Ltd., Watford, Herts. WD18 7JJ, UK. Text revised October 2012.

MEDIJEL[®] Gel

Lidocaine hydrochloride

Aminoacridine hydrochloride

Oral Gel For Common Mouth Ulcers

***FAST effective relief
from common mouth ulcers,
sore gums & denture-rubbing***

MEDIJEL[®] Gel

Lidocaine hydrochloride

Aminoacridine hydrochloride

Oral Gel For Common Mouth Ulcers

***FAST effective relief
from common mouth ulcers,
sore gums & denture-rubbing***

If symptoms persist for more than 7 days, consult your doctor or dentist. Do not use Medijel Gel if you are allergic to Lidocaine or any of the other ingredients.

Instructions for use: The local anaesthetic action of Medijel Gel provides fast relief from common mouth ulcers, sore gums and denture rubbing. Apply a small quantity of Medijel Gel on a clean fingertip directly to the painful area. Repeat the application every 20 minutes if necessary. Use only as directed. Do not store above 25°C.



5 011784 030126 >