

Olbas[®]

**blackcurrant
pastilles**

Olbas[®] **blackcurrant pastilles**

Manufactured by
Ernest Jackson & Co. Ltd.,
CREDITON, DEVON, EX17 3AP, UK,
for the Marketing
Authorisation holder
G. R. Lane Health Products Ltd.,
Sisson Road, Gloucester,
GL2 0GR, UK, to whom any letters
should be addressed.



Lanes
HEALTH

40g^e

Before you take Olbas Blackcurrant Pastilles

Do not take if you are allergic to any of the ingredients.

Taking other medicines Please tell your doctor or pharmacist if you are taking or have recently taken any other medicines, including medicines obtained without a prescription.

Pregnancy and breast-feeding It is best not to take any non-essential medication when pregnant or breast-feeding. Ask your doctor or pharmacist for advice before taking any medicine.

Important information about some of the ingredients of Olbas Blackcurrant Pastilles Contains a source of phenylalanine. May be harmful for people with phenylketonuria. If you have been told by your doctor that you have an intolerance to Maltitol, contact your doctor before taking this medicinal product.

How to take Olbas Blackcurrant Pastilles

Adults and children aged 16 and over: Dissolve 1 pastille slowly in the mouth when required. Do not take more than 8 pastilles in any 24 hour period. Not recommended for children under 16 years old. You must contact a doctor if symptoms worsen or do not improve after 7 days.

If you take more Olbas Blackcurrant Pastilles than you should, talk to a doctor or pharmacist immediately.

Possible side effects

Like all medicines, Olbas Blackcurrant Pastilles can cause side effects, although not everybody gets them. Tell your doctor if you notice any of the following mild side effects: Heartburn, wheezing, worsening of asthma or shortness of breath. If any of the side effects become serious, or if you notice any side effect not listed on this carton, please tell your doctor or pharmacist.

How to store Olbas Blackcurrant Pastilles

Keep out of the reach and sight of children.

Do not use Olbas Blackcurrant Pastilles after the expiry date which is stated on the carton.

Do not store above 25°C.

Do not use if sealed inner pouch is cut or torn.