



Vicks Sinex Soother 0.5mg/ml nasal spray solution Oxymetazoline Hydrochloride

Patient information leaflet

In this leaflet:

1. What Vicks Sinex Soother is and what it is used for
2. Before you use Vicks Sinex Soother
3. How to use Vicks Sinex Soother
4. Possible side effects
5. How to store Vicks Sinex Soother
6. Further information

Read all of this leaflet carefully because it contains important information for you.

This medicine is available without prescription. However, you still need to use Vicks Sinex Soother carefully to get the best results from it.

- **Keep this leaflet.** You may need to read it again.
- Ask your pharmacist if you need more information or if you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4.
- You should contact a doctor if your symptoms worsen or do not improve after 3 days.

1. What Vicks Sinex Soother is and what it is used for

Vicks Sinex Soother is a nasal spray used for localised relief of the symptoms of a blocked nose in association with a cold or rhinitis. The effect of the spray starts after several minutes and lasts for 6-8 hours.

2. Before you use Vicks Sinex Soother

Do not use Vicks Sinex Soother if you:

- are allergic (hypersensitive) to the active substance oxymetazoline hydrochloride or any of the other ingredients of Vicks Sinex Soother (see section 6)
- are taking (or have taken during the last 2 weeks) monoamine oxidase inhibitors (MAO inhibitors are used in the treatment of Parkinson's disease and depression)
- suffer from acute heart disease or cardiac asthma
- suffer from increased eyeball-pressure (narrow-angle glaucoma)
- have had an operation to remove your pituitary gland

- have inflamed skin or mucous membranes of your nostrils or have scabs in your nose.

Do not give Vicks Sinex Soother to children under 12 years of age

Take special care with Vicks Sinex Soother

Before using ask your doctor for advice if you have:

- high blood pressure
- heart disease, including angina
- diabetes
- thyroid problems
- enlarged prostate gland (prostatic hypertrophy)

Taking other medicines

Please inform your doctor or pharmacist before using Vicks Sinex Soother if you are taking, or have recently taken, any other medicines, even those obtained without a prescription.

- **monoamine oxidase inhibitors (MAOIs)** are used in the treatment of Parkinson's disease and depression. **Do not use** Vicks Sinex Soother if you have taken MAOIs during the last two weeks.
- **tricyclic antidepressants** - inform your doctor if you are taking drugs used in the treatment of depression such as amitriptyline and imipramine.

Beta blockers or high blood pressure medicines

– inform your doctor before use if you are taking any of these.

Parkinson's disease medicine (bromocriptine)

– inform your doctor before use if you are taking this medicine.

Pregnancy and Breast-feeding

Ask your doctor or pharmacist for advice before using Vicks Sinex Soother. The spray should be used with caution during pregnancy or if breast feeding.

Driving and using machines

No effect on ability to drive or use machines has been observed.

Important information about some of the ingredients of Vicks Sinex Soother

Benzalkonium chloride (a preservative) is an ingredient in Vicks Sinex Soother which might cause swelling of nasal mucosa, especially during long-term use.

3. How to use Vicks Sinex Soother

Always use Vicks Sinex Soother exactly as your pharmacist or doctor has told you. You should check with your doctor or pharmacist if you are not sure.

► **Dosage:** The recommended dose is

Adults and children over 12 years: 1-2 sprays in each nostril no more than 2-3 times daily.

Children under 12 years - Do not use.

Do not exceed the recommended dose.

Do not use for longer than 5-7 days in a row. If used over a longer period, Vicks Sinex Soother may cause a blocked nose.

Contact your doctor if symptoms worsen or do not improve after 3 days.

► **Administration:**

1. Remove the cap. Keep the bottle upright with the nasal applicator just into the nostril. Hold the other nostril closed.
2. Spray and breathe in deeply through your nose at the same time. Repeat for the other nostril.
3. Adults and children over 12 years may repeat steps 1 and 2 once if necessary.
4. Wipe the nasal applicator with a clean, wet tissue and replace the cap immediately after application.

► **If you use more Vicks Sinex Soother than you should**

If you use too much Vicks Sinex Soother and you experience any unwanted effects or symptoms, seek medical advice immediately and take the bottle or pack with you.

► **If you forget to use Vicks Sinex Soother**

Do not take a double dose if you forget to take the previous dose; instead, continue using according to the administration instructions.

If you have any further questions on the use of this product, ask your pharmacist.

4. Possible side effects

Like all medicines, Vicks Sinex Soother can cause side effects, although not everybody gets them.

Uncommon (occurring in less than 1 in 100 people): Sneezing, dryness or irritation in the nose, mouth or throat.

Rare (occurring in less than 1 in 1000 people): Anxiety, fatigue, irritability, disturbed sleep in children, rapid heartbeat, palpitations, raised blood pressure, blocked nose, swelling of the nasal lining, headache, nausea, flushing, rash and visual disturbances.

► **Reporting of side effects**

If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via the Yellow Card Scheme at:
www.mhra.gov.uk/yellowcard
By reporting side effects you can help provide more information on the safety of this medicine.

5. How to store Vicks Sinex Soother

Keep out of the reach and sight of children. Do not store above 25°C.

Do not use Vicks Sinex Soother after the expiry date which is printed on the carton. The expiry date refers to the last day of the month stated.

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

6. Further information

What Vicks Sinex Soother contains

The **active substance** is: Oxymetazoline hydrochloride 0.5 mg/ml.

The **other ingredients** are: sorbitol, sodium citrate dihydrate, tyloxapol, chlorhexidine digluconate solution, citric acid anhydrous, aloe dry extract, benzalkonium chloride solution, acesulfame potassium, menthol (levo), cineole, disodium edetate (dihydrate), carvone (levo), sodium hydroxide, purified water. (See end of section 2 for benzalkonium chloride).

What Vicks Sinex Soother looks like and contents of the pack

Vicks Sinex Soother is a metered dose nasal pump spray. The pack contains an amber glass bottle with a white pump on top. The bottle is filled with 10 ml or 15 ml of product.

Not all pack sizes may be marketed.

Marketing Authorisation Holder and Manufacturer

Marketing Authorisation Holder:
Procter & Gamble(Health & Beauty Care) Limited,
Weybridge, Surrey KT13 OXP United Kingdom

Manufacturer:

Procter & Gamble, Manufacturing GmbH
Procter & Gamble Strasse 1, 64521 Gross-Gerau,
Germany